



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

the earliest birds of that species, transients in our latitude, pass hurriedly, and stop only in such kinds of country as most resemble their ultimate destination. In other words, as a general rule, their occurrence in the State in spring becomes extended from the higher lands to the lower as the season advances.

---

## THE WINTER RANGES OF THE WARBLERS (MNIOTILTIDÆ).

BY W. W. COOKE.

MANY expert ornithologists have spent a great deal of time and care in working out the breeding range of each species of North American birds. No comparable effort has been bestowed on the question of the winter home, and as a result many loose statements are current in ornithological literature. There seems to be a tendency to consider any record south of the United States as a wintering record, whereas quite a number of species pass in migration through the West Indies or Central America to winter in South America.

The following tables show at a glance what part of the district south of the United States is occupied as a winter home and what is crossed in migration. It might be added that the charts are not designed to show anything with reference to the breeding range of any of these species; so that the statement, for instance, that *Protonotaria citrea* occurs in migration in the southeastern United States is not to be taken as indicating that it does not breed in that same district.

The tables are condensed from the Bulletin on the 'Distribution and Migration of North American Warblers' recently issued by the Biological Survey.



		S. E. United States	Bahamas	Cuba	Jamaica	Haiti	Porto Rico	Lesser Antilles	Trinidad	Texas	S. W. United States	Mexico	Yucatan	Guatemala	Honduras	Nicaragua	Costa Rica	Panama	Colombia	Venezuela	Guitana	Brazil	Ecuador	Pern	Bolivia	Chile
32	<i>Dendroica blackburniae</i>	+	+									+	+	+	+				+				+			
33	<i>d. dominica</i>	+	+									+	+	+	+											
34	<i>d. albiflora</i>									+			+													
35	<i>graciae</i>									+			+													
36	<i>nigrescens</i>																									
37	<i>chrysoparia</i>																									
38	<i>virens</i>																									
39	<i>townsendi</i>																									
40	<i>occidentalis</i>																									
41	<i>kirtlandi</i>																									
42	<i>vigorsii</i>																									
43	<i>palmarum</i>																									
44	<i>p. hypochrysea</i>																									
45	<i>discolor</i>																									
46	<i>Seiurus aurocapillus</i>																									
47	<i>noveboracensis</i>																									
48	<i>n. notabilis</i>																									
49	<i>motacilla</i>																									
50	<i>Geothlypis formosa</i>																									
51	<i>agilis</i>																									
52	<i>philadelphia</i>																									
53	<i>tolmiei</i>																									
54	<i>trichas</i>																									
55	<i>t. occidentalis</i>																									
56	<i>t. ignota</i>																									
57	<i>t. arizela</i>																									
58	<i>t. brachidactyla</i>																									
59	<i>t. sinuosa</i>																									
60	<i>beltingi</i>																									
61	<i>poliocephala</i>																									
62	<i>Icteria virens</i>																									
63	<i>v. longicauda</i>																									
64	<i>Wilsonia mitrata</i>																									

— = Occurs in migration

+ = Winters

\* = Rare or accidental

[illegible]